

Treatment-Emergent Integrase Strand Transfer Inhibitor (INSTI) Resistance-Associated Mutations Among People With HIV-1 Treated With Dolutegravir (DTG) + Lamivudine (3TC) With Pre-existing M184V/I From Real-world and Interventional Studies

Dainielle Fox,¹ Gary Blick,² Thibault Mesplède,³ Gustavo Verdier,⁴ Mónica Calderón,¹ Chris M. Parry,⁵ Richard Grove,⁶ Emilio Letang,⁷ Julie Priest,¹ Bryn Jones,⁵ Gregory Huhn^{1*} ¹ViiV Healthcare, Durham, NC, USA; ²Health Care Advocates International, Stratford, CT, USA; ³Erasmus University Medical Center, Rotterdam, Netherlands; ⁴ViiV Healthcare, Montréal, Canada; ⁵ViiV Healthcare, London, UK; ⁶GSK, London, UK; ⁷ViiV Healthcare, Tres Cantos, Madrid, Spain

*Presenting on behalf of the authors.

Key Takeaways

A systematic literature review (SLR) identified 300 publications representing 47,350 unique people using dolutegravir (DTG) + lamivudine (3TC) in both real-world and interventional settings

Treatment-emergent integrase strand transfer inhibitor (INSTI) resistance rarely (<0.05%) developed in people initiating or switching to DTG + 3TC

None of the people who developed treatment-emergent INSTI resistance had known pre-existing

These data support the in vitro findings that indicate that M184V/I antagonizes the development of resistance against DTG

Introduction

- DTG + 3TC has a high barrier to resistance and is recommended as first-line and maintenance antiretroviral therapy (ART) for people with HIV-1¹⁻³
- In vitro, tissue cultures exposed to DTG monotherapy selected INSTI resistance (R263K) with wild-type virus but not with M184V/I-containing virus⁴
- Though the M184V/I resistance mutation confers high-level resistance to 3TC, data suggest that M184V/I presence does not increase the risk of developing INSTI resistance at virologic failure (VF) in people with HIV-1 using DTG + 3TC
- Archived M184V/I detected in proviral DNA did not impact efficacy in people with virologic suppression switching to DTG/3TC in randomized phase 3 trials (TANGO, n=4; SALSA, n=5)^{5,6}
- People who switched to DTG + 3TC while virologically suppressed had low rates of VF and no cases of treatmentemergent INSTI resistance, regardless of historical or current M184V/I, in the VOLVER study at Week 48,⁷ the SOLAR-3D study through Week 144,⁸ and in a meta-analysis of 12 real-world and interventional studies⁹
- Outcomes data, especially real-world evidence (RWE), continue to emerge for people with M184V/I using DTG + 3TC
- To assess the latest evidence on the impact of M184V/I on treatment-emergent INSTI resistance, we present an updated SLR with studies reporting DTG + 3TC use through March 2024 and include populations naive to ART

Methods

Systematic Literature Review

- An SLR was conducted following Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines
- Ovid MEDLINE[®], Embase[®], PubMed, Cochrane library databases, and relevant congresses were searched for RWE studies reporting DTG + 3TC use in people with HIV-1 published between January 2013 and March 2024, as described previously⁹
- Phase 3/3b DTG + 3TC randomized controlled trials from ViiV Healthcare were added to the interventional study pool post hoc
- Any records excluded from RWE study screening for having an interventional study design were screened separately for analysis inclusion

Analysis

- Descriptive analyses were independently performed for 2 groups of SLR-identified studies with different screening criteria:
- Studies evaluating M184V/I at baseline: included all publications with ≥10 people using DTG + 3TC that evaluated baseline M184V/I (even if 0 people with baseline M184V/I are reported)
- Studies in which M184V/I was an exclusion criterion were excluded, apart from 2 interventional studies that reported archived resistance data^{6,7}
- For people naive to ART, baseline is the time of DTG + 3TC initiation; for people with prior ART use, baseline is the time of switch to DTG + 3TC
- All studies: included all publications with ≥10 people using DTG + 3TC
- After screening, "lead" studies with the most data and/or highest n per cohort were identified and used to represent unique populations within each group and avoid double-counting people across multiple and/or overlapping publications or overlapping with another cohort
- Lead studies among studies evaluating M184V/I at baseline may include non-lead studies from the group of all studies
- Analysis denominators use the total n of people using DTG + 3TC, not the total n of people with resistance tests performed

Acknowledgments: This study was funded by ViiV Healthcare. Editorial assistance and graphic design support for this poster were provided under the direction of the authors by Fingerpaint Medical (formerly MedThink SciCom) and funded by ViiV Healthcare. Data included in this poster have previously been presented in full at HIV Drug Therapy Glasgow; November 10-13, 2024; Glasgow, Scotland; Poster P330.

Results

Included Studies and Populations

- The SLR identified 300 publications (n=249 RWE, n=51 interventional) that reported ≥10 people with HIV-1 using DTG + 3TC • These studies represented 108 discrete cohorts and trials and 47,350 unique people with HIV-1 after accounting for population overlap
- The number of lead studies included for each analysis group is summarized in the Table
- 30 lead studies (N=10,383) were included in the analysis group of studies evaluating M184V/I at baseline (n=1049 naive to ART; n=9318 prior ART use; n=16 not specified)
- 108 lead studies (N=47,350) were included in the analysis group of all studies, regardless of whether M184V/I was evaluated at baseline (n=3404 naive to ART; n=21,912 prior ART use; n=22,034 not specified)

Figure 1. PRISMA Flowchart



^aPublications excluded due to interventional study design were screened separately for inclusion in the analysis. ^bSearched congresses included the Asian Conference on Hepatitis and AIDS; Australasian HIV & AIDS Conference; British Association for Sexual Health and HIV; British HIV Association; Canadian Conference on HIV/AIDS Research; Conference on Retroviruses and Opportunistic Infections; European AIDS Clinical Society; Grupo de Estudio del SIDA-SEIMC; HIV & Hepatitis in the Americas; HIV Glasgow; IDWeek; International AIDS Society/International Conference; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on Antiviral Research; International Conference; International Conference; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on Antiviral Research; International Conference; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Transmitted Infections in Africa; International Conference on AIDS and Sexually Trans Francophone Conference on HIV/Hepatitis/Sexual Health/Emerging Infections (AFRAVIH; searched May 24, 2022); Japanese Society for AIDS Research; Kenya Association of Physicians; National Conference of AIDS Society of India; Société Française de Lutte contre le Sida; and The HIV Netherlands Australia Thailand Research Collaboration. Includes all studies with M184V/I at baseline, even n=0, and excludes studies with M184V/I as an exclusion criterion. regardless of whether M184V/I at baseline is evaluated (ie, includes studies with M184V/I presence as an exclusion criterion and where M184V/I may be present but not reported)

Prevalence of Baseline M184V/I

- Overall prevalence of M184V/I at baseline was 5% (512/10,383) among studies evaluating M184V/I at baseline and 0.9% (406/47,350) among all studies (Table)
- 4 of the cohorts that were included in both analysis groups were each represented by a different lead study between analysis groups, due to differences in screening criteria
- Thus, the analysis of all studies has a higher total n than the analysis of studies evaluating M184V/I at baseline, which provides a higher estimate of people with baseline M184V/I
- Prevalence of M184V/I reported by interventional studies was 20% (219/1096) and 8% (221/2914) among studies evaluating M184V/I at baseline and among all studies, respectively (Table)
- Several of these interventional studies focus on M184V/I and thus overestimate its prevalence

Table. Summary of SLR-Identified Publications Reporting People Using DTG + 3TC by Study Type

	Lead ^a RWE studies	Lead ^a interventional studies	Total
Studies evaluating M184V/I at baseline ^b			
Studies, n	22	8	30
People using DTG + 3TC, n	9287	1096	10,383
People with baseline M184V/I, n (%)	293 (3) ^c	219 (20)	512 (5)
RNA or unspecified	165 (2) ^c	201 (18)	366 (4)
Proviral DNA	128 (1) ^d	18 (2)	146 (1)
All studies ^e			
Studies, n	84	24	108
People using DTG + 3TC, n	44,436	2914	47,350
People with baseline M184V/I, n (%)	185 (0.4) ^c	221 (8)	406 (0.9)
RNA or unspecified	162 (0.4) ^c	202 (7)	364 (0.8)
Proviral DNA	23 (0.05) ^d	19 (0.7)	42 (0.1)
^a The publication with the largest n for each RWE cohort or interventional trial was selected after all SLR-identified publications were screened for inclusion. ^b Include exclusion criterion. ^c 1 study ¹⁰ reported that n=60 people overall had historical M18 at least n=15 people using DTG + 3TC had historical M184V/I. ^d n=2 confirmed threw with M184V/I presence as an exclusion criterion and where M184V/I may be preserved.	selected to represent that cohort/trial, to account s studies that report any number of individuals 4V/I (DTG + 3TC, n=174; DTG + RPV, n=46) a ough personal communications. ^e Includes all st ent but not reported).	nt for potentially overlapping populations. Lead studies for ea with M184V/I at baseline, even n=0, and excludes studies wi nd at least n=1 person using DTG + RPV did not have histor udies regardless of whether M184V/I at baseline is evaluate	ach analysis group were ith M184V/I as an rical M184V/I; therefore, id (ie, includes studies
Treatment-Emergent INSTI Resistance			
• Figure 2 summarizes INSTI resistance detected	at VF in each group of studie	S:	
 2/10,383 (0.02%) people (n=512 with known pre- 	existing M184V/I) among studie	es evaluating M184V/I at baseline	
 6/47,350 (0.01%) people (n=406 with known pre- 	existing M184V/I) among all stu	udies	
 None of the people who developed treatment-en 	nergent INSTI resistance had	known pre-existing M184V/I	
Figure 2. Treatment-Emergent INSTI Resistance or Lead Interventional Studies (A) Evaluating M	Reported at VF With DTG + 184V/I at Baseline and (B) C	- 3TC Across Populations From Lea Overall	ad RWE
A Studies evoluating M404)/// at he	B B		



alncludes studies that report any number of individuals with M184V/I at baseline, even n=0, and excludes studies with M184V/I as an exclusion criterion. Includes all studies regardless of whether M184V/I at baseline, even n=0, and excludes studies with M184V/I as an exclusion criterion. baseline is evaluated (ie, includes studies with M184V/I presence as an exclusion criterion and where M184V/I may be present but not reported).

Conclusions

- with DTG + 3TC, as was previously reported for in vitro tissue culture studies⁴
- Overall rates of VF were very low across all types of studies included in this analysis
- increase the risk of INSTI mutations developing at VF
- histories are unavailable, or in cases where M184V/I is inadvertently missed

References: 1. EACS. https://www.eacsociety.org/media/guidelines-12.0.pdf. Accessed September 30, 2024. 2. Gandhi et al. JAMA. 2023;329:63-84. 3. DHHS. https://clinicalinfo.hiv.gov/sites/default/files/guidelines/documents/adult-adolescent-arv/guidelines-adult-adolescent-arv.pdf. Accessed September 30, 2024. 4. Oliveira et al. A/DS. 2016;30:2267-2273. 5. Wang et al. CROI 2020; Boston, MA. Poster 489. 6. Underwood et al. CROI 2022; Virtual. Poster 481. 7. De Miguel et al. EACS 2023; Warsaw, Poland. Poster eP.A.097. 8. Blick et al. AIDS 2024; Munich, Germany. Oral presentation OAX0903LB. 9. Kabra et al. Open Forum Infect Dis. 2023;10:ofad526. 10. Galizzi et al. Int J Antimicrob Agents. 2020;55:105893.

• No people with HIV-1 with known pre-existing M184V/I experienced treatment-emergent INSTI resistance development at VF • One potential explanation for these findings is that M184V/I may have a protective effect against the development of INSTI resistance

• This SLR further supports the growing body of evidence showing that use of DTG + 3TC in the presence of M184V/I does not

• This provides reassurance for considering DTG/3TC as a first-line or switch option if resistance test results or complete treatment